

SHOULDER ARTHROSCOPY WITH POSTERIOR LABRAL REPAIR REHABILITATION PROTOCOL

General Notes

As tolerated should be understood to include with safety for the surgical procedure; a sudden increase in pain, swelling, or other undesirable factors are indicators that you are doing too much too soon. If any of these occur, decrease activity level and ice.

During rehabilitation if there are any neurovascular findings please call the office.

Ice should be applied to the shoulder for 15-20 minutes following each exercise, therapy, or training session. Return to sport is based on provider team (physician, physician assistant, athletic trainer, therapist) input and appropriate testing.

All times and exercises are to serve as guidelines. Progression through the protocol should be based upon criteria as opposed to dates listed and will vary depending on each individual patient. Progress should be agreed upon by the patient and his/her team of providers.

Post-Operative Phase I: Healing Phase - (Day 1 - Week 4)

Goals:

- Minimize shoulder pain and inflammatory response
- Protect the integrity of the surgical repair
- Gradually restore pain free passive range of motion (PROM)
- Enhance/ensure adequate scapular function

Sling:

- To be worn at all times except when bathing or performing therapy

PROM:

- Glenohumeral (GH) flexion to 90 degrees
- Abduction in the plane of the scapula to 90 degrees
- Internal rotation (IR) to 30 degrees at 30 degrees of abduction
- External rotation (ER) to 60 degrees at 30-40 degrees of abduction; respect anterior capsule tissue integrity with ER ROM.

SHOULDER ARTHROSCOPY WITH POSTERIOR LABRAL REPAIR REHABILITATION PROTOCOL

Therapeutic Exercises:

- Pendulum exercises
- Scapular stabilization exercises
- Ball squeezes
- Active assisted and active range of motion (AAROM/AROM: wrist, fingers, and supination/pronation with arm in sling (at 90 degrees of elbow flexion)
- Active elbow flexion/extension
- At week 3: Initiate Submaximal GH isometrics: flexion, extension, abduction, IR, ER

Modalities/Education:

- Frequent cryotherapy for pain and inflammation
- Pre-modulated electrical stimulation to shoulder for pain reduction
- Patient education regarding posture, joint protection, positioning, hygiene, etc.

Manual:

- Scar and soft tissue mobilization as needed
- Joint mobilizations as needed

Post-Operative Phase II: Motion Phase - (Week 4 – Week 6)

Goals:

- Minimize shoulder pain and inflammatory response
- Protect the integrity of the surgical repair
- Progress ROM
- Begin light waist level activities

Sling:

- Discontinue use of pillow as directed by physician after week 4.
- Begin to wean from sling between weeks 5-6 as directed by MD.
- Discontinue sling as directed by physician after week 6

ROM:

- Progress shoulder ROM (do not force any painful motion)
 - Forward flexion to 120 degrees
 - Abduction in the plane of the scapula to 100 degrees
 - IR to 45 degrees at multiple angles of abduction
 - ER to 75 degrees at 0-40 degrees of abduction then progress to multiple angles of abduction

SHOULDER ARTHROSCOPY WITH POSTERIOR LABRAL REPAIR REHABILITATION PROTOCOL

Therapeutic Exercises:

- Gentle Posterior capsular stretching as needed
- Progress to AAROM/AROM exercises of the shoulder with proper GH rhythm
- Continue AROM of elbow, wrist, and hand
- Progress scapular stabilization exercises

Modalities:

- Continue cyrotherapy for pain and inflammation
- Pre-modulated electrical stimulation to shoulder for pain reduction

Manual:

- Scar and soft tissue mobilization as needed
- Joint mobilizations as needed

Post-Operative Phase III: Strengthening Phase (Week 6 – Week 8)

Goals:

- Achieve normal GH and scapular rhythm
- Progress A/PROM
- Remain pain free for all exercises

ROM

- Progress shoulder ROM (do not force any painful motion)
 - Forward flexion to 140 degrees
 - Abduction in the plane of the scapula to 140 degrees
 - IR to 75 degrees at multiple angles of abduction
 - ER to WNL

Therapeutic Exercises:

- Progress as tolerated, concentric and eccentric exercises
 - Achieve PROM/AROM in a given plane before strengthening in that plane
- Initiate:
 - Closed chain activities
 - UBE: no resistance
 - Maximal Isometrics and gentle isotonic with arm at side
 - Scapular Stabilization with gentle resistance

SHOULDER ARTHROSCOPY WITH POSTERIOR LABRAL REPAIR REHABILITATION PROTOCOL

Manual:

- Scar and soft tissue mobilizations as needed
- Joint mobilizations as needed

Cardio:

- Stationary bike, elliptical with light upper body, stair climber

Modalities:

- Continue cyotherapy for pain and inflammation
- Pre-modulated electrical stimulation to shoulder for pain reduction
- Continued patient education: posture, joint protection, positioning, hygiene, etc.

Post-Operative Phase IV: Activity Phase - (Week 10 – Week 16)

Goals:

- Progress strength, endurance, neuromuscular control
- Begin shoulder height functional activities
- Gradual and planned buildup of stress to anterior joint capsule

ROM

- Progress shoulder ROM to WNL (do not force any painful motion)

Therapeutic Exercises:

- Rotator Cuff Strengthening - ER/IR strengthening
 - Side lying ER with towel roll
 - Manual resistance to ER in supine in scapular plane
 - ER/IR with exercise tubing at 0 degrees of abduction (towel roll)
- Prone rowing at 30/45/90 degrees of abduction to neutral arm position
- Begin rhythmic stabilization drills at week 8
 - ER/IR in the scapular plane
 - Flexion/extension and adduction/abduction at various angles of elevation
- Progress AROM to isotonic: flexion, scaption, abduction
- Continue A/PROM as needed
- Progress strengthening to focus on both upper and lower segments:
 - PNF patterns
 - IR resistive band at 45, 90 degrees of abduction
- Increase resistance for UBE
- Progress isotonic strengthening if patient demonstrates no compensatory strategies, is not painful, and has no residual soreness

SHOULDER ARTHROSCOPY WITH POSTERIOR LABRAL REPAIR REHABILITATION PROTOCOL

Precautions:

- Do not overstress the posterior capsule with aggressive overhead activities/strengthening
- Avoid contact sports/activities
- Patient education regarding a gradual increase to shoulder activities

Manual:

As needed

Post-Operative Phase V: Return to Sport Phase (Week 16+)

Goals:

- Maintain full non-painful P/AROM
- Progress to full strenuous work, throwing, and overhead activities

Precautions:

- Avoid excessive posterior capsule stress
- Do not begin plyometrics, throwing, or overhead athletic moves until 4 months post-op or cleared by MD.

Therapeutic Exercise:

- Continue to advance scapula and upper extremity strengthening as tolerated in all planes
 - Include: diagonal/functional patterns, 90/90 strengthening, and CKC exercises

Cardio:

- Stationary bike, jogging/running/sprinting on treadmill, elliptical, rowing, kick board in swimming pool, stair climber

Milestones for return to sport activities and clearance:

- Completion and passing of shoulder functional test at MD PT clinic
- No complaints of pain or instability
- Adequate ROM for task completion bilaterally
- Full strength and endurance of rotator cuff and scapular musculature for task completion bilaterally
- Regular completion of home exercise program